AUTHOR INDEX

Bold type indicates authors of commentaries, editorials, and Resource Corners; (No.) indicates issue number.

Abraham WT, 82
ACCENT I Study Group, 92
Alcorn J, 117
Alderson P, 51
Allen G, 84
Alper BS, 52
American Lung Association Asthma Clinical Research Centers, 18
Anderson CS, 114
Anderson I, 101
Andrews SJ, 112
Antithrombotic Trialists' Collaboration, 5
Arozullah AM,
Arroll B, 116, 109
Asplund K, 5
Assendelft WJ, 65
Atlantic Cardiovascular Patient Outcomes Research Team (C-PORT), 43
Aversano LT, 43
Aversano T, 43

Baime MJ, 25
Bainton CE, A-16 (No. 1)
Ballew KA, 46
Barnes PJ, 19
Barry M, 61
Bates ER, 2
Baum KD, A-14 (No. 1)
Beck JC, 57
Becker LA, 72, 16
Beiser A, 76
Bell-Syer SE, 97
Belluzi A, 91
Bender B, 37
Bennett SG, 37
Bernal W, 117
Berry E, 110
Binnie R, 107
Birks J, 10
Bisson JI, 69
Blackhouse G, 91
Blecker B, 86–87
Blecker SB, 86–87
Boersma E, 2
Bond CM, 15
Booth A, A-14 (No. 2)
Boothroyd D, 105
Borghi L, 62
Borzak S, 1
Boulvain M, 30
Bradley CP, 34
Brazzelli M, 28
Bridges-Webb C, 106
Brezis M, 94
Broadhurst RJ, 114
Brooks PT, 112
Brophy MT, 47
Bucher HC, 49
Buerkle G, 44
Buettner HJ, 44
Bulpitt CJ, 95
Bunn F, 51
Burr M, 63
Califf RM, 85
Cameron C, 90
Cannon CP, 1
Carruthers SG, 83
Cascino GD, 75
Cates C, 104
Ceftin and Flonase for Sinusitis (CAFFS) Investigators, 14
Chalmers B, 68
Chan BK, 42
Chan FK, 13
Chessman A, 24, 113
Chick J, 27
Chin A, 109
Choi P, 51
Christakis DA, 96
Coates AL, 107
Combination Hemotherapy and Mortality Prevention (CHAMP) Study Group, 47
Cooke J, 95
Cordes JA, 74
Corbett EC, 90
Couriel JM, 64
Cramer JA, 27
Crawford F, 97
Curhan G, 62

D'Souza AL, 95
Dahlén SE, 20
Dahlöf B, 86–87
Daley J, 34
Davidson I, 33
Davidson J, 69
Dawson DR, 22
de Benedictis FM,
Del Mar CB, A-14 (No. 3), 116
Demers C, 48
Dempster C, 101
Dennis CL, 68
Devereux RB, 86
Diabetes Prevention Program Research Group, 55
Dick J, 36
DiGiulio GA, 31
Dixon A, 73
Dixon ME, 26
Dolor RJ, 14
Donaldson N, 117
Douketis J, 42
Dubé C, 54
Ducharme FM, 32, 64
Duda SH, 12

Ebbert JO, A-15 (No. 1)
Egger M, 57
Elley CR, 109
Elliott G, 111
Entwistle V, 36
Epstein AE, 82
Erkinjuntti T, 102
Ernst E, 25
EuroSIDA Study Group, 35
Evans JG, 10
Ezekowitz MD, 47

Fahey T, 106
Fanelli CG, 89
Fay JK, 63
Fazel S, 76
Feagan BG, 92, 115
Fekete T, 112
Ferrucci L, 4
Feudtner C, 96
Finney JW, 58
Fiore LD, 47
Fisher C, 110
Fisher WG, 82
Fleming MF, 59
Flynn CA, 72
Forbes S, 114
Fraser DG, 33
Freemantle N, 101
French MT, 59
Furberg CD, 4

Gallop R, 68
Ganda O, 3
Gard P, 16
Garrison MM, 96
Gatell JM, 35
Gauthier S, 102
Gaynes BN, 100
Geba GP,
Gijtenbeek JM, 66
Gillespie LD, 9
Gillespie WJ, 9
Glanville J, 101
Glasziou P, 90
Glazier RH, 72
Gluckman R, 35, 99
Gluud C, 17
Goeree R, 91
Goldberg J, 70
Gonzalez-del-Rey JA, 31
Goodman SG, 85
Gøtzsche PC, 93, 98
Grad R, 23
Graham A, 58–59
Gray J, 4
Green RJ, 32
Greenstone M, 63
Greenwood R, 22
Greyber E, 88
Griffiths P, 28
Grimshaw JM 15,
Grosser D, 19, 71

Hall J, 81
Hall PF, 67
Hammer A, 57
Hamoud M, 8
Hanauer SB, 92
Hanbury DC, 112
Hankey GJ, 114
Harper D, 15
Harrington RA, 2, 43

Harris AD, 95
Hart R, 97
Hayden M, 6
Haynes J, 105
He J, 109
Helkamp AS, 14
Henderson WG, 34
Hengstler P, 49
Herkner H, 24
HERS Research Group, 105
Hesdorffer DC, 75
Heslin J, 22
Hiatt WR, 12
Hill K, 9
Hillegass WE, 82
Hirsch C, 57
Hlatky MA, 81, 105
Hodnett E, 29, 68
Hofstee DJ, 66
Holmes W, 16
Holroyd-Leduc J, A-15 (No. 2)
Hoogland PH, 66
Huang ES, 3
Hull RD, 11
Hulscher ME, 99
Hypothermia after Cardiac Arrest Study Group, 46

Ibsen H, 87
Iliffe S, 23
International Study Group, 32
Ioannou GN, 94
IONA Study Group, 83
Ishani A, 61

Jain R, 6
Jamrozik K, 84, 114
Jewell D, 67
Johansen HK, 98
Johnson SC, 77
Jones A, 63
Jong P, 48

Kearon C, 33
Kelder JC, 45
Kelly S, 110
Kenealy T, 116
Kenny RA, 8
Kenyon S, 30
Kerry S, 56
Kerse N, 116
Khan KS, 30
Khuri SF, 34
King PD, 60
Kirk GF, 27
Kitzman DW, 48
Kjaergard LL, 17
Kjeldsen SE, 86
Korula J, 93
Kowdley K, 94
Kozak E, 7
Kreder HS, 65
Krogsgaard K, 17
Krol WF, 27
Krystal JH, 27
Kudva YC, 56, 89
Kurz A, 102
Kuster GM, A-16 (No. 1)
Kwan J, 77

Lancaster T, 52
Lang AE, 53
Lang W, 24
Langhorne P, 114
Larson E, 102
Lefebvre C, 51
Leibovici L, 94
Lesko SM, 108
Lewin SA, 36
Lewis PR, 52
Lichtenstein GR, 92
LIFE study group, 86–87
Lindholm LH, 87
Liu PP, 48
Loder E, 88
Loftus EV Jr, 115
Logan RF, 92
Logroscino G, 75
Londborg P, 69
Louik C, 108
Lowe J, 41
Ludbrook A, 15
Lundgren JD, 35
Luria JW, 31
Luxenberg JS, 10
Luz O, 12

Mac Donald R, 61
Macfarlane J, 16
Malenka DJ, 44
Malmström K, 20
Marshall JC, 36
Marshall JK, 91
Martel AL, 33
Massel D, 45
Martin W, 53
MASTER Anaesthesia Trial Study Group, 84
Mattock M, 56
Mazur LJ, 108
McGinn T, A-11–12 (No. 2)
McKelvie RS, 48
McQuay HJ, 103, 52
Meier G, 49
Meigs JB, 3
Menza MA, 21
Meschi T, 62
Meyerhoff J, 26
Miller J, 42
Minder CE, 57
Minster A, 113
MIRACLE Study Group, 82
Mitchell AA, 108
Miyasaki JM, 53
Mocroft A, 35
Moffatt ME, 28
Moliterno DJ, 2
Møller AM, 7
Mollison J, 15
Montori VM, A-15 (No. 1), 55
Moody AR, 33
Moore A, 103
Moran JL, 50
Morgan PS, 33
Morrow DA, 1
Morton SC, 99
Moss AJ, 81
Moyer A, 58
Mueller C, 44
Mulrow C, 6
Multicenter Automatic Defibrillator Implantation Trial II Investigators, 81
Mundt MP, 59
Myers RP, 60
Myles PS, 84

Neilson J, 30
Nelson HD, 42
Newby LK, 85
Niewoehoefer D, 20
Nikodem C, 29
Nixon J, A-14 (No. 1)
Nizankowska E, 20
Noël PH, 74

O'Byrne PM, 19
O'Donnell M, 11
O'Malley PG, 74
Oh VM, 47
Oldman AD, 103
Ormiston T, 104
Oudkerk M, 111

Pampanelli S, 89
Paulseth JE, 53
Participants in the National Veterans Affairs Surgical Quality Improvement Program, 34
Passamani E, 43
Pasternak G, 88
Paton JY, 31
Pearlstein T, 69
Pedersen T, 7
Penninx BW, 4
Peter JV, 50
Petty DR, 73
Pfisterer M, A-16 (No. 1)
Phillips C, 6
Phillips-Hughes J, 50
Pickup J, 56
Pignone M, 6, 74, 100
Pineo GF, 11
Pirozzo S, 90
Plokker HW, 45
Polis AB, 26
Poole P, 104
Porcellati F, 89
Powell J, 22
Poynard T, 60
Rajkumar C, 95

Ramirez G, 74
Ramsey SD, A-11–12 (No. 3)
Raynor DK, 73
Regimbeau C, 60
Reichen J, 17
Richardson DA, 8
Richardson WS, A-11–12 (No. 1)
Rifai N, 1
Rigg JR, 84
Robertson MC, 9
Rodriguez-Roisin R, 19
Roose SP, 21
Rosen RC, 21
Rosenberg M, 6
Rosenheck RA, 27
Rostasy K, 113
Rotstein C, 98
Rowe BH, 71, 18
Rushton JL, 100
Rutks I, 61

Salpeter E, 88, 104
Salpeter S, 88, 104
Sandborn WJ, 115
Sandercock P, 77
Saunders LD, 71
Schianchi T, 62
Schindler C, 49
Schnitzer TJ, 26
Schoenfeld P, 115
Schroeder K, 106
Schuh S, 107
Schwartz M, 54
Seidman SN,
Selhub J, 76
Seshadri S, 76
Shabsigh R, 21
Shafran S, 97
Sheehan B, 76
Shannon KC, 37
Shepperd S, 23
Sherman SE, 100
Shiau JY, 54
Shukla VK, 54
Sigal RJ, 88
Simon GE, 101
Simon JA, 49
Simmer K, 68
Sinacore JM, 37
Singer DE, 3
Sinuff T, 50
Skea ZC, 36
Skelly MM, 92
Smidt N, 65
Smith AL, 82
Smith B, 19, 71
Smith D, 101
Smith LA, 103
Smucny JJ,
Soares-Weiser K, 94
Spooner CH, 71
Stafstrom CE, 113
Stanbrook ME, 104
Stark G, 61
Steen N, 8
Stein PD, 11
Stevens CA, 64
Stone EG, 99
Stuck AE, 57
Suchowersky O, 53
Summerbell C, 90
Suttorp MJ, 45
Swearingen CE, 58

TACTICS-TIMI 18 Investigators, 1
ten Berg JM, 45
Tepe G, 12
Teper A, 32
Thoennissen J, 24
TIME Investigators, A-16 (No. 1)
To KF, 13
Tønnesen H, 7
Tourette's Syndrome Study Group, 70
Tur-Kaspa R, 94

Uhari M, 96

VACT Group, 26
van Beek EJ, 111
van der Windt DA, 65
Van Niel CW, 96
Veldhuyzen van Zanten SJO, 13
Vergun P, 58
Verheugt FW, 45
Veterans Affairs Naltrexone Cooperative Study 425 Group, 27
Vezina RM, 108
Villebro N, 7
Vittinghoff E, 105

Walker JM, A-16 (No. 1)
Warn D, 50
Watson MC, 15
Weaver AL, 26
Weiner WJ, 53
Wendon J, 117
Wesseldine LJ, 64
Westwood ME, 110
Whelton SP, 109
White HD, 85
Whitehead A, 10
Wiebe S, 75
Wiedemann H, 72
Wielopolski P, 111
Wiener SL, 66
Wilcock GK, 10
Willet LR, 14
Williams JW Jr, 74
Wilson SA, 21
Wilt T, 61
Witsell DL, 14
Women's Health Initiative Investigators, 41
Wu JC, 13
Wyer PC, 107
Wyncoll D, 117

Xin X, 109

Young G, 67

Zareba W, 81
Zermansky AG, 73
Zwarenstein M, 36

A

Abdomen
- Epidural anesthesia and analgesia did not reduce most comorbid outcomes in high-risk patients having major abdominal surgery, 84

Accidental falls
- Review: Intrinsic and environmental risk-factor modification reduces falls in elderly persons, 9

Acetaminophen
- Arterial blood lactate measurements quickly identified risk for death from paracetamol-induced liver failure, 117
- Ibuprofen was more protective against asthma morbidity than acetaminophen in asthmatic children with fever, 108
- Rofecoxib, 25 mg/d, was more effective than rofecoxib, 12.5 mg/d, celecoxib, or acetaminophen in osteoarthritis of the knee, 26

Acetates
- Montelukast moderately decreased asthma symptoms in children with persistent asthma (Correction), A-17 (No. 1)

Acidosis, lactic
- Review: Metformin does not increase fatal or nonfatal lactic acidosis or blood lactate levels in type 2 diabetes mellitus, 89

Adrenergic beta-agonists
- Review: β2-agonists are ineffective but increase adverse effects in acute bronchitis without underlying pulmonary disease, 72

Adrenergic beta-agonists
- Review: Cardioselective β-blockers did not reduce respiratory function in patients with chronic obstructive pulmonary disease, 92

Alcohol drinking
- Review: Brief interventions reduce drinking in patients not seeking treatment, 58

Alcoholism
- A brief intervention reduced alcohol drinking for up to 48 months in problem drinkers, 59
- Naltrexone for 3 or 12 months did not reduce drinking in alcohol dependence, 27

Alzheimer disease
- Galantamine improved cognition and global functioning in vascular dementia or Alzheimer disease with cerebrovascular disease, 102
- Review: Selegiline leads to a small short-term improvement in cognition and activities of daily living in Alzheimer disease, 10

Analgesia, epidural
- Epidural anesthesia and analgesia did not reduce most comorbid outcomes in high-risk patients having major abdominal surgery, 84

Androstadienes
- Growth rate was greater with fluticasone propionate than with beclomethasone dipropionate in children with chronic asthma, 32

Anesthesia, general
- Epidural anesthesia and analgesia did not reduce most comorbid outcomes in high-risk patients having major abdominal surgery, 84

Angina
- An early invasive strategy was effective in the acute coronary syndrome with elevated cardiac troponin I and T levels, 1
- Nicorandil reduced coronary events in stable angina, 83
- Review: Glycoprotein IIb/IIIa inhibitors reduced death or MI in acute coronary syndromes not routinely scheduled for revascularization, 2

Angiography
- Magnetic resonance angiography had high specificity but moderate sensitivity for detecting pulmonary emboli, 111

Angioplasty
- Isotonic hydration reduced contrast media–associated nephropathy in elective or emergency coronary angioplasty, 44
- Primary percutaneous intervention was more effective than thrombolytic therapy for acute myocardial infarction, 43
- Revascularization led to less angina and fewer adverse cardiac events than did optimal medical care in angina pectoris in the elderly (Letter), A-16 (No. 1)

Angiotensin-converting enzyme inhibitors
- Review: Angiotensin-receptor blockers do not reduce mortality or hospitalization rates in heart failure, 48

Antiasthmatic agents
- Montelukast moderately decreased asthma symptoms in children with persistent asthma (Correction), A-17 (No. 1)

Antibiotics
- Delayed prescription reduced antibiotic use in the common cold, 116
- Review: Antibiotic prophylaxis reduces mortality and bacterial infection in cirrhosis and gastrointestinal bleeding, 94
- Review: Antibiotics improve maternal and fetal outcomes and are safe in preterm, prelabor rupture of membranes, 30
- Review: Probiotics are effective in preventing antibiotic-associated diarrhea, 95
- Verbal advice plus an information leaflet reduced antibiotic use in acute bronchitis, 16

Antibodies, monoclonal
- Maintenance infliximab increased time to loss of response in active Crohn disease, 92

Anticholesteremic agents
- Review: Interventions that lower cholesterol levels or blood pressure in diabetic patients prevent cardiovascular disease, 3

Antidepressive agents
- Review: Venlafaxine is more effective than selective serotonin-reuptake inhibitors for depression, 101

Antiemetics
- Review: Antiemetic drugs reduce nausea in early pregnancy, 67

Antifungal agents
- Review: Antifungal agents do not reduce mortality in neutropenia caused by chemotherapy or bone marrow transplantation, 98
- Review: Oral and intravaginal agents are equally effective for treatment of uncomplicated vulvovaginal candidiasis, 15
- Review: Oral antifungal drugs promote cure of fungal foot infections, 97
- Antihypertensive agents in diabetic patients prevent cardiovascular disease, 3
- Treating isolated systolic hypertension prevented major cardiovascular events across strata of risk in older patients, 4

Anti-inflammatory agents, non-steroidal
- *Helicobacter pylori* eradication reduced the risk for ulcers in dyspeptic patients who were about to begin NSAID treatment, 13
- Rofecoxib, 25 mg/d, was more effective than rofecoxib, 12.5 mg/d, celecoxib, or acetaminophen in osteoarthritis of the knee, 26
- Oral dexamethasone improved respiratory distress in children < 2 years of age with acute bronchiolitis, 107

Antiretroviral therapy
- A simple scoring system predicted clinical progression in HIV patients receiving highly active antiretroviral therapy, 35

Antitussive agents
- Review: Lack of evidence exists for effectiveness of over-the-counter cough preparations for children with URTI, 106

Arterial occlusive diseases
Abciximab added to urokinase increased amputation-free survival in peripheral arterial occlusion of the legs, 12
Arthroplasty, replacement, hip
Review: Extended out-of-hospital prophylaxis with heparin prevents deep venous thrombosis in elective hip arthroplasty, 11
Aspirin
Review: Aspirin reduces CAD events in persons with no history of cardiovascular disease, but it increases gastrointestinal bleeding (Correction), 80
Warfarin combined with low-dose aspirin in MI did not provide clinical benefit beyond that of aspirin alone, 47
Asthma
Ibuprofen was more protective against asthma morbidity than acetaminophen in asthmatic children with fever, 108
Low-dose budesonide improved asthma control in mild asthma; adding formoterol improved control in corticosteroid-treated patients, 19
Montelukast improved pulmonary function and asthma-specific quality of life in aspirin-intolerant asthma, 20
Montelukast moderately decreased asthma symptoms in children with persistent asthma (Correction), A-17 (No. 1)
Parental education and guided self-management of asthma or wheeze in preschool children was not effective, 64
Asthma, exercise-induced
Review: Nedocromil sodium decreases airflow reduction and recovery time in exercise-induced bronchoconstriction, 71
Atenolol
Losartan reduced cardiovascular morbidity and mortality more than atenolol in patients with diabetes and essential hypertension, 87
Losartan reduced strokes and new-onset diabetes more than atenolol in essential hypertension, 86
Treating isolated systolic hypertension prevented major cardiovascular events across strata of risk in older patients, 4
Attention deficit disorder with hyperactivity
Clonidine and methylphenidate were effective for attention deficit hyperactivity disorder in children with comorbid tics, 70

B

Beclomethasone
Growth rate was greater with fluticasone propionate than with beclomethasone dipropionate in children with chronic asthma, 32
Review: Inhaled corticosteroids for asthma or chronic obstructive pulmonary disease do not increase bone loss, 63
Bed rest
Bed rest, physiotherapy, and continuation of activities of daily living were equally effective for sciatica, 66
Behavior therapy
Review: Behavioral interventions plus laxatives are effective for defecation disorders, but biofeedback does not add benefit, 28
Review: Brief interventions reduce drinking in patients not seeking treatment, 58
Birthing centers
Review: A home-like birth environment has beneficial effects on labor and delivery, 29
Blood pressure
Review: Aerobic exercise reduces systolic and diastolic blood pressure in adults, 109
Body height
Growth rate was greater with fluticasone propionate than with beclomethasone dipropionate in children with chronic asthma, 32
Bone density
Review: Inhaled corticosteroids for asthma or chronic obstructive pulmonary disease do not increase bone loss, 63
Brain diseases
Mild hypothermia improved neurologic outcome and reduced mortality after cardiac arrest because of ventricular arrhythmia, 46
Brain injury
A multidisciplinary community-based rehabilitation program improved social functioning in severe traumatic brain injury, 22
Breast feeding
Telephone-based peer support increased the duration of breast-feeding in primiparous mothers, 68
Bronchiolitis
Oral dexamethasone improved respiratory distress in children < 2 years of age with acute bronchiolitis, 107
Bronchitis
Review: β2-agonists are ineffective but increase adverse effects in acute bronchitis without underlying pulmonary disease, 72
Verbal advice plus an information leaflet reduced antibiotic use in acute bronchitis, 16
Bronchodilator agents
Low-dose budesonide improved asthma control in mild asthma; adding formoterol improved control in corticosteroid-treated patients, 19
Budesonide
Low-dose budesonide improved asthma control in mild asthma; adding formoterol improved control in corticosteroid-treated patients, 19
Review: Inhaled corticosteroids for asthma or chronic obstructive pulmonary disease do not increase bone loss, 63

C

Calcium, dietary
A normal-calcium, low-protein, low-salt diet reduced recurrence of renal stones over 5 years, 62
Candidiasis vulvovaginal
Review: Oral and intravaginal agents are equally effective for treatment of uncomplicated vulvovaginal candidiasis, 15
Cardiac arrest
Mild hypothermia improved neurologic outcome and reduced mortality after cardiac arrest because of ventricular arrhythmia, 46
Cardiac pacing, artificial
Cardiac pacing reduced nonaccidental falls in older adults with cardioinhibitory carotid sinus hypersensitivity, 8
Cardiac resynchronization was effective for moderate-to-severe heart failure with intraventricular conduction delay, 82
Carotid stenosis
Review: Magnetic resonance angiography is accurate for detecting high-grade carotid artery stenosis that is suitable for surgery, 110
Cefuroxime
Adding intranasal fluticasone to cefuroxime resolved acute rhinosinusitis, 14
Cerebrovascular disorders
Galantamine improved cognition and global functioning in vascular dementia or Alzheimer disease with cerebrovascular disease, 102
Predictors of poor outcome in acute stroke included dependence before stroke, severe hemiparesis, and recurrent stroke, 114

Review: Antiplatelet therapy prevents occlusive vascular events in high-risk patients, 5
Review: In-hospital care pathways for acute stroke do not improve clinical outcomes but do lower quality of life, 77
Cervical vertebrae
Review: Bed rest does not prevent cervical or lumbar puncture headaches, 24
Chlorthalidone
Treating isolated systolic hypertension prevented major cardiovascular events across strata of risk in older patients, 4
Clonidine
Clonidine and methylphenidate were effective for attention deficit hyperactivity disorder in children with comorbid tics, 70
Colonic neoplasms
Incorporating economic analysis into clinical practice guidelines: a guide for users, A-11–12 (No. 3)
Common cold
Delayed prescription reduced antibiotic use in the common cold, 116
Coronary angiography
Isotonic hydration reduced contrast media–associated nephropathy in elective or emergency coronary angioplasty, 44
Planned angiography after percutaneous coronary intervention increased reinterventions but did not affect mortality or MI, 45
Coronary disease
Estrogen plus progestin increased coronary heart disease and breast cancer events in postmenopausal women, 41
Planned angiography after percutaneous coronary intervention increased reinterventions but did not affect mortality or MI, 45
Revascularization led to less angina and fewer adverse cardiac events than did optimal medical care in angina pectoris in the elderly (Letter), A-16 (No. 1)
Review: Aspirin reduces CAD events in persons with no history of cardiovascular disease, but it increases gastrointestinal bleeding (Correction), 80
Review: N-3 polyunsaturated fatty acids reduce fatal MI, sudden death, and overall mortality in coronary artery disease, 49
Review: Oral glycoprotein IIb/IIIa inhibitors increase mortality and myocardial infarction, 85
Costs and cost analysis
Incorporating economic analysis into clinical practice guidelines: a guide for users, A-11–12 (No. 3)
Cough
Review: Lack of evidence exists for effectiveness of over-the-counter cough preparations for children with URTI, 106
Review: β2-agonists are ineffective but increase adverse effects in acute bronchitis without underlying pulmonary disease, 72
Critical illness
Review: Albumin increases mortality in critically ill patients, 51
Critical pathways
Review: In-hospital care pathways for acute stroke do not improve clinical outcomes but do lower quality of life, 77
Crohn disease
Maintenance infliximab increased time to loss of response in active Crohn disease, 92
Review: Infliximab increases response and remission rates in fistulizing or treatment-resistant Crohn disease, 91
Review: One tenth of patients with Crohn disease have prolonged remission, 115
Croup
Oral dexamethasone led to fewer treatment failures than did nebulized dexamethasone or placebo in children with mild croup, 31
Cyclooxygenase inhibitors
Rofecoxib, 25 mg/d, was more effective than rofecoxib, 12.5 mg/d, celecoxib, or acetaminophen in osteoarthritis of the knee, 26

D

Decision making
The Resourceful Patient (Resource Corner), A-14 (No. 3)
Defibrillators, implantable
A prophylactically implanted cardioverter defibrillator reduced all-cause mortality in myocardial infarction, 81
Delivery of health care
A comprehensive annotated reminder tool increased appropriate screening in primary care, 37
Delivery rooms
Review: A home-like birth environment has beneficial effects on labor and delivery, 29
Dementia
Elevated plasma total homocysteine levels increased the risk for dementia in the elderly, 76
Galantamine improved cognition and global functioning in vascular dementia or Alzheimer disease with cerebrovascular disease, 102
Depression
1 of 2 quality-improvement interventions for depression in managed care was more effective but more costly than usual care (Correction), A-17 (No. 1)
Review: Questionnaires for detecting clinical depression in primary care have similar diagnostic accuracy, 74
Review: Screening for depression reduces persistent depression, 100
Review: Venlafaxine is more effective than selective serotonin-reuptake inhibitors for depression, 101
Sildenafil improved erectile dysfunction and quality of life in men with comorbid mild-to-moderate depression, 21
Dexamethasone
Oral dexamethasone improved respiratory distress in children < 2 years of age with acute bronchiolitis, 107
Oral dexamethasone led to fewer treatment failures than did nebulized dexamethasone or placebo in children with mild croup, 31
Diabetes mellitus
Irbesartan reduced progression of nephropathy caused by type 2 diabetes independent of the effect on blood pressure (Correction), A-17 (No. 1)
Irbesartan was renoprotective in patients with type 2 diabetes, hypertension, and microalbuminuria (Correction), A-17 (No. 1)
A lifestyle intervention or metformin prevented or delayed the onset of type 2 diabetes in persons at risk, 55
Overall satisfaction increased more with inhaled insulin than with subcutaneous insulin in type 1 diabetes (Correction), A-17 (No. 1)
Review: Intensive insulin therapy with a subcutaneous infusion pump slightly improves glycemic control in type 1 diabetes, 56
Review: Interventions that lower cholesterol levels or blood pressure in diabetic patients prevent cardiovascular disease, 3
Review: Metformin does not increase fatal or nonfatal lactic acidosis or blood lactate levels in type 2 diabetes mellitus, 88
A split regimen of regular insulin at dinner and NPH insulin at bedtime reduced nocturnal hypoglycemia in type 1 diabetes, 89

Diabetic nephropathies
Irbesartan reduced progression of nephropathy caused by type 2 diabetes independent of the effect on blood pressure (Correction), A-17 (No. 1)
Irbesartan was renoprotective in patients with type 2 diabetes, hypertension, and microalbuminuria (Correction), A-17 (No. 1)
Diarrhea
Review: *Lactobacillus* is safe and effective for treating children with acute infectious diarrhea, 96
Review: Probiotics are effective in preventing antibiotic-associated diarrhea, 95
Diet
A normal-calcium, low-protein, low-salt diet reduced recurrence of renal stones over 5 years, 62
Review: Advice on low-fat diets is not better than other weight-reducing diets for sustaining weight loss in obesity, 90
Dopamine agonists
Review: Selegiline improves symptoms and levodopa is better than pramipexole for motor function in untreated Parkinson disease, 53
Drug hypersensitivity
Montelukast improved pulmonary function and asthma-specific quality of life in aspirin-intolerant asthma, 20
Dyspepsia
Helicobacter pylori eradication reduced the risk for ulcers in dyspeptic patients who were about to begin NSAID treatment, 13
Review: Proton-pump inhibitors alleviate symptoms of functional (nonulcer) dyspepsia but may not be better than H2-antagonists, 54

E

Echinacea
Review: St. John's wort, ginkgo, saw palmetto, and kava may be effective for some conditions, 25
Esophageal and gastric varices
Review: Somatostatin analogues reduce blood products used in acute bleeding esophageal varices but do not reduce mortality, 93
Estrogen replacement therapy
Effects of hormone therapy on quality of life in postmenopausal women with CAD differed according to menopausal symptoms, 105
Estrogen plus progestin increased coronary heart disease and breast cancer events in postmenopausal women, 41
Review: Postmenopausal estrogen replacement therapy increases the risk for venous thromboembolism, 42
Ethanolamines
Low-dose budesonide improved asthma control in mild asthma; adding formoterol improved control in corticosteroid-treated patients, 19
Evidence-based medicine
Critical Evaluation of Research in Physical Rehabilitation: Towards Evidence-Based Practice (Resource Corner), A-15 (No. 2)
Textbook descriptions of disease—where's the beef? (Editorial), A-11 (No. 1)
The Resourceful Patient (Resource Corner), A-14 (No. 3)
TRIP database (Resource Corner), A-15 (No. 2)
Exercise
Cardiac resynchronization was effective for moderate-to-severe heart failure with intraventricular conduction delay
Review: Aerobic exercise reduces systolic and diastolic blood pressure in adults, 109

F

Family practice
1 of 2 quality-improvement interventions for depression in managed care was more effective but more costly than usual care (Correction), A-17 (No. 1)
Fatty acids, omega-3
Review: N-3 polyunsaturated fatty acids reduce fatal MI, sudden death, and overall mortality in coronary artery disease, 49
Fecal incontinence
Review: Behavioral interventions plus laxatives are effective for defecation disorders, but biofeedback does not add benefit, 28
Fetal membranes, premature rupture
Review: Antibiotics improve maternal and fetal outcomes and are safe in preterm, prelabor rupture of membranes, 30
Fever
Ibuprofen was more protective against asthma morbidity than acetaminophen in asthmatic children with fever, 108
Fibrinolytic agents
Review: Aspirin reduces CAD events in persons with no history of cardiovascular disease, but it increases gastrointestinal bleeding (Correction), 80
Fluid therapy
Isotonic hydration reduced contrast media–associated nephropathy in elective or emergency coronary angioplasty, 44
Review: Albumin increases mortality in critically ill patients, 51
Fluticasone
Review: Inhaled corticosteroids for asthma or chronic obstructive pulmonary disease do not increase bone loss, 63
Foot dermatoses
Review: Oral antifungal drugs promote cure of fungal foot infections, 97
Fractures
Review: Inhaled corticosteroids for asthma or chronic obstructive pulmonary disease do not increase bone loss, 63

G

Galantamine
Galantamine improved cognition and global functioning in vascular dementia or Alzheimer disease with cerebrovascular disease, 102
Gastroenteritis
Review: *Lactobacillus* is safe and effective for treating children with acute infectious diarrhea, 96
Gastrointestinal hemorrhage
Review: Antibiotic prophylaxis reduces mortality and bacterial infection in cirrhosis and gastrointestinal bleeding, 94
Review: Somatostatin analogues reduce blood products used in acute bleeding esophageal varices but do not reduce mortality, 93
Geriatric assessment
Cardiac pacing reduced nonaccidental falls in older adults with cardioinhibitory carotid sinus hypersensitivity, 8
Ginkgo biloba
Review: St. John's wort, ginkgo, saw palmetto, and kava may be effective for some conditions, 25
Glucocorticoids, synthetic
Adding intranasal fluticasone to cefuroxime resolved acute rhinosinusitis, 14
Growth rate was greater with fluticasone propionate than with beclomethasone dipropionate in children with chronic asthma, 32

Physiotherapy or a wait-and-see policy were the best options for lateral epicondylitis at 1 year, 65
Guideline adherence
A comprehensive annotated reminder tool increased appropriate screening in primary care, 37

H

Headache
Children's drawings of headache pain were accurate for diagnosing migraine, 113
Review: Bed rest does not prevent cervical or lumbar puncture headaches, 24
Health education
Parental education and guided self-management of asthma or wheeze in preschool children was not effective, 64
Health services for the aged
Review: Home visiting with multidimensional assessment and multiple visits is effective in low-risk elderly persons, 57
Heart failure, congestive
Cardiac resynchronization was effective for moderate-to-severe heart failure with intraventricular conduction delay, 82
Review: Angiotensin-receptor blockers do not reduce mortality or hospitalization rates in heart failure, 48
Helicobacter pylori
Helicobacter pylori eradication reduced the risk for ulcers in dyspeptic patients who were about to begin NSAID treatment, 13
Heparin, low-molecular-weight
Review: Extended out-of-hospital prophylaxis with heparin prevents deep venous thrombosis in elective hip arthroplasty, 11
Hepatitis C
Review: Interferon α-2b is effective for biochemical and virologic outcomes in acute hepatitis C virus infection, 60
Herpes zoster
Review: Tricyclic antidepressants, capsaicin, gabapentin, and oxycodone are effective for postherpetic neuralgia, 52
HIV infections
A simple scoring system predicted clinical progression in HIV patients receiving highly active antiretroviral therapy, 35
Home care services
A multidisciplinary community-based rehabilitation program improved social functioning in severe traumatic brain injury, 22
Review: Hospital-at-home care for early discharge or admission avoidance does not improve health outcomes, 23
Homocysteine
Elevated plasma total homocysteine levels increased the risk for dementia in the elderly, 76
Hypertension
Irbesartan reduced progression of nephropathy caused by type 2 diabetes independent of the effect on blood pressure (Correction), A-17 (No. 1)
Irbesartan was renoprotective in patients with type 2 diabetes, hypertension, and microalbuminuria (Correction), A-17 (No. 1)
Losartan reduced cardiovascular morbidity and mortality more than atenolol in patients with diabetes and essential hypertension, 87
Losartan reduced strokes and new-onset diabetes more than atenolol in essential hypertension, 86
Hypoglycemia
A split regimen of regular insulin at dinner and NPH insulin at bedtime reduced nocturnal hypoglycemia in type 1 diabetes, 89
Hypoglycemic agents
Review: Intensive insulin therapy with a subcutaneous infusion pump slightly improves glycemic control in type 1 diabetes, 56
Review: Interventions that lower cholesterol levels or blood pressure in diabetic patients prevent cardiovascular disease, 3
Hypothermia, induced
Mild hypothermia improved neurologic outcome and reduced mortality after cardiac arrest because of ventricular arrhythmia, 46

I

Ibuprofen
Ibuprofen was more protective against asthma morbidity than acetaminophen in asthmatic children with fever, 108
Impotence
Sildenafil improved erectile dysfunction and quality of life in men with comorbid mild-to-moderate depression, 21
Infliximab
Maintenance infliximab increased time to loss of response in active Crohn disease, 92
Review: Infliximab increases response and remission rates in fistulizing or treatment-resistant Crohn disease, 91
Influenza vaccine
The inactivated trivalent split-virus influenza vaccine is safe in stable asthma, 18
Institutionalization
Predictors of poor outcome in acute stroke included dependence before stroke, severe hemiparesis, and recurrent stroke, 114
Insulin infusion systems
Review: Intensive insulin therapy with a subcutaneous infusion pump slightly improves glycemic control in type 1 diabetes, 56
Insulin
A split regimen of regular insulin at dinner and NPH insulin at bedtime reduced nocturnal hypoglycemia in type 1 diabetes, 89
Overall satisfaction increased more with inhaled insulin than with subcutaneous insulin in type 1 diabetes (Correction), A-17 (No. 1)
Review: Intensive insulin therapy with a subcutaneous infusion pump slightly improves glycemic control in type 1 diabetes, 56
Interferon
Review: Interferon a-2b is effective for biochemical and virologic outcomes in acute hepatitis C virus infection, 60
Review: Interferon-a plus ribavirin improves the virologic response in chronic hepatitis C, 17
Internet
TRIP database (Resource Corner), A-15 (No. 2)

K

Kava
Review: St. John's wort, ginkgo, saw palmetto, and kava may be effective for some conditions, 25
Kidney calculi
A normal-calcium, low-protein, low-salt diet reduced recurrence of renal stones over 5 years, 62

L

Lactates
Arterial blood lactate measurements quickly identified risk for death from paracetamol-induced liver failure, 117
Lactobacillus
Review: *Lactobacillus* is safe and effective for treating children with acute infectious diarrhea, 96

Leukotriene antagonists
Montelukast improved pulmonary function and asthma-specific quality of life in aspirin-intolerant asthma, 20
Levodopa
Review: Selegiline improves symptoms and levodopa is better than pramipexole for motor function in untreated Parkinson disease, 53
Lifestyle
A lifestyle intervention or metformin prevented or delayed the onset of type 2 diabetes in persons at risk, 55
Liver cirrhosis
Review: Antibiotic prophylaxis reduces mortality and bacterial infection in cirrhosis and gastrointestinal bleeding, 94
Liver failure, acute
Arterial blood lactate measurements quickly identified risk for death from paracetamol-induced liver failure, 117
Losartan
Irbesartan reduced progression of nephropathy caused by type 2 diabetes independent of the effect on blood pressure (Correction), A-17 (No. 1)
Losartan reduced cardiovascular morbidity and mortality more than atenolol in patients with diabetes and essential hypertension, 87
Losartan reduced strokes and new-onset diabetes more than atenolol in essential hypertension, 86

M

Magnetic resonance angiography
Magnetic resonance angiography had high specificity but moderate sensitivity for detecting pulmonary emboli, 111
Review: Magnetic resonance angiography is accurate for detecting high-grade carotid artery stenosis that is suitable for surgery, 110
Magnetic resonance imaging
Magnetic resonance direct thrombus imaging was accurate in suspected deep venous thrombosis, 33
Mass screening
Incorporating economic analysis into clinical practice guidelines: a guide for users, A-11–12 (No. 3)
Review: Organizational change and patient involvement may increase the use of prevention cancer screening services, 99
Review: Screening for depression reduces persistent depression, 100
MEDLINE
MEDLINEplus: A golden gateway to health information resources (Resource Corner), A-14 (No. 2)
Metformin
A lifestyle intervention or metformin prevented or delayed the onset of type 2 diabetes in persons at risk, 55
Review: Metformin does not increase fatal or nonfatal lactic acidosis or blood lactate levels in type 2 diabetes mellitus, 88
Methylphenidate
Clonidine and methylphenidate were effective for attention deficit hyperactivity disorder in children with comorbid tics, 70
Migraine
Children's drawings of headache pain were accurate for diagnosing migraine, 113
Review: Several drugs, especially triptans, are effective for pain relief in acute migraine, 103
Monitoring, physiologic
Early goal-direct therapy reduced mortality and multiorgan dysfunction in severe sepsis or septic shock (Correction), 80
Myocardial infarction
Primary percutaneous intervention was more effective than thrombolytic therapy for acute myocardial infarction, 43
A prophylactically implanted cardioverter defibrillator reduced all-cause mortality in myocardial infarction, 81
Review: Antiplatelet therapy prevents occlusive vascular events in high-risk patients, 5
Review: Glycoprotein IIb/IIIa inhibitors reduced death or MI in acute coronary syndromes not routinely scheduled for revascularization, 2
Review: Oral glycoprotein IIb/IIIa inhibitors increase mortality and myocardial infarction, 85
Warfarin combined with low-dose aspirin in MI did not provide clinical benefit beyond that of aspirin alone, 47
Myocardial revascularization
Planned angiography after percutaneous coronary intervention increased reinterventions but did not affect mortality or MI, 45
Revascularization led to less angina and fewer adverse cardiac events than did optimal medical care in angina pectoris in the elderly (Letter), A-16 (No. 1)

N

Naltrexone
Naltrexone for 3 or 12 months did not reduce drinking in alcohol dependence, 27
Nausea
Review: Antiemetic drugs reduce nausea in early pregnancy, 67
Nebulizers and vaporizers
Oral dexamethasone led to fewer treatment failures than did nebulized dexamethasone or placebo in children with mild croup, 31
Nedocromil
Review: Nedocromil sodium decreases airflow reduction and recovery time in exercise-induced bronchoconstriction, 71
Neoplasms
Review: Antifungal agents do not reduce mortality in neutropenia caused by chemotherapy or bone marrow transplantation, 98
Review: Organizational change and patient involvement may increase the use of prevention cancer screening services, 99
Neuralgia
Review: Tricyclic antidepressants, capsaicin, gabapentin, and oxycodone are effective for postherpetic neuralgia, 52
Neutropenia
Review: Antifungal agents do not reduce mortality in neutropenia caused by chemotherapy or bone marrow transplantation, 98
Nicorandil
Nicorandil reduced coronary events in stable angina, 83

O

Obesity
Review: Advice on low-fat diets is not better than other weight-reducing diets for sustaining weight loss in obesity, 90
Osteoarthritis, knee
Rofecoxib, 25 mg/d, was more effective than rofecoxib, 12.5 mg/d, celecoxib, or acetaminophen in osteoarthritis of the knee, 26

P

Pain measurement
Rofecoxib, 25 mg/d, was more effective than rofecoxib, 12.5 mg/d, celecoxib, or acetaminophen in osteoarthritis of the knee, 26

Parkinson disease
Review: Selegiline improves symptoms and levodopa is better than pramipexole for motor function in untreated Parkinson disease, 53
Patient education
MEDLINEplus: A golden gateway to health information resources (Resource Corner), A-14 (No. 2)
Review: Intrinsic and environmental risk-factor modification reduces falls in elderly persons, 9
Verbal advice plus an information leaflet reduced antibiotic use in acute bronchitis, 16
Patient participation
Pharmacists changed more repeat prescriptions for elderly patients than did usual general practice review, 73
Patient satisfaction
Overall satisfaction increased more with inhaled insulin than with subcutaneous insulin in type 1 diabetes (Correction), A-17 (No. 1)
Patient-centered care
Review: Interventions for health care providers improve provider-patient interactions and patient satisfaction, 36
Peptic ulcer
Helicobacter pylori eradication reduced the risk for ulcers in dyspeptic patients who were about to begin NSAID treatment, 13
Pharmacists
Pharmacists changed more repeat prescriptions for elderly patients than did usual general practice review, 73
Physical therapy techniques
Bed rest, physiotherapy, and continuation of activities of daily living were equally effective for sciatica, 66
Critical Evaluation of Research in Physical Rehabilitation: Towards Evidence-Based Practice (Resource Corner), A-15 (No. 2)
Physiotherapy or a wait-and-see policy were the best options for lateral epicondylitis at 1 year, 65
Physician's practice patterns
Pharmacists changed more repeat prescriptions for elderly patients than did usual general practice review, 73
Phytotherapy
Review: *Pygeum africanum* extracts improve symptoms and urodynamics in symptomatic benign prostatic hyperplasia, 61
Piperazines
Sildenafil improved erectile dysfunction and quality of life in men with comorbid mild-to-moderate depression, 21
Platelet aggregation inhibitors
Abciximab added to urokinase increased amputation-free survival in peripheral arterial occlusion of the legs, 12
Review: Antiplatelet therapy prevents occlusive vascular events in high-risk patients, 5
Platelet glycoprotein GPIIb-IIIa complex
Review: Oral glycoprotein IIb/IIIa inhibitors increase mortality and myocardial infarction, 85
Review: Glycoprotein IIb/IIIa inhibitors reduced death or MI in acute coronary syndromes not routinely scheduled for revascularization, 2
Pneumonia
A risk index with 14 variables predicted 30-day postoperative pneumonia after major noncardiac surgery, 34
Postmenopause
Effects of hormone therapy on quality of life in postmenopausal women with CAD differed according to menopausal symptoms, 105
Estrogen plus progestin increased coronary heart disease and breast cancer events in postmenopausal women, 41
Review: Postmenopausal estrogen replacement therapy increases the risk for venous thromboembolism, 42
Postoperative complications
A preoperative smoking intervention decreased postoperative complications in elective knee or hip replacement, 7
A risk index with 14 variables predicted 30-day postoperative pneumonia after major noncardiac surgery, 34
Pregnancy
Review: Antibiotics improve maternal and fetal outcomes and are safe in preterm, prelabor rupture of membranes, 30
Review: Antiemetic drugs reduce nausea in early pregnancy, 67
Preoperative care
A preoperative smoking intervention decreased postoperative complications in elective knee or hip replacement, 7
Preventive health services
A comprehensive annotated reminder tool increased appropriate screening in primary care, 37
Review: Home visiting with multidimensional assessment and multiple visits is effective in low-risk elderly persons, 57
Review: Organizational change and patient involvement may increase the use of prevention cancer screening services, 99
Primary health care
Review: Interventions for health care providers improve provider-patient interactions and patient satisfaction, 36
Review: Screening for depression reduces persistent depression, 100
Verbal advice plus an information leaflet reduced antibiotic use in acute bronchitis, 16
Probiotics
Review: *Lactobacillus* is safe and effective for treating children with acute infectious diarrhea, 96
Review: Probiotics are effective in preventing antibiotic-associated diarrhea, 95
Prostatic hyperplasia
Review: *Pygeum africanum* extracts improve symptoms and urodynamics in symptomatic benign prostatic hyperplasia, 61
Proton pumps
Review: Proton-pump inhibitors alleviate symptoms of functional (nonulcer) dyspepsia but may not be better than H_2-antagonists, 54
Psychotherapy, brief
A brief intervention reduced alcohol drinking for up to 48 months in problem drinkers, 59
Pulmonary disease, chronic obstructive
Review: Cardioselective b-blockers did not reduce respiratory function in patients with chronic obstructive pulmonary disease, 92
Pulmonary embolism
Magnetic resonance angiography had high specificity but moderate sensitivity for detecting pulmonary emboli, 111

Q

Questionnaires
Review: Questionnaires for detecting clinical depression in primary care have similar diagnostic accuracy, 74

R

Reading
Users' Guides to the Medical Literature Interactive Web site (Resource Corner), A-14 (No. 1)

Receptors, angiotensin
Irbesartan was renoprotective in patients with type 2 diabetes, hypertension, and microalbuminuria (Correction), A-17 (No. 1)
Review: Angiotensin-receptor blockers do not reduce mortality or hospitalization rates in heart failure, 48
Rehabilitation
Critical Evaluation of Research in Physical Rehabilitation: Towards Evidence-Based Practice (Resource Corner), A-15 (No. 2)
A multidisciplinary community-based rehabilitation program improved social functioning in severe traumatic brain injury, 22
Respiratory insufficiency
Review: Noninvasive ventilation reduces mortality in acute respiratory failure, 50
Respiratory tract infections
Review: Lack of evidence exists for effectiveness of over-the-counter cough preparations for children with URTI, 106
Review: A home-like birth environment has beneficial effects on labor and delivery, 29
Review: N-3 polyunsaturated fatty acids reduce fatal MI, sudden death, and overall mortality in coronary artery disease, 49
Rhinitis
Adding intranasal fluticasone to cefuroxime resolved acute rhinosinusitis, 14
Ribavirin
Review: Interferon-α plus ribavirin improves the virologic response in chronic hepatitis C, 17
Risk assessment
A risk index with 14 variables predicted 30-day postoperative pneumonia after major noncardiac surgery, 34

S

Sciatica
Bed rest, physiotherapy, and continuation of activities of daily living were equally effective for sciatica, 66
Selegiline
Review: Selegiline improves symptoms and levodopa is better than pramipexole for motor function in untreated Parkinson disease, 53
Review: Selegiline leads to a small short-term improvement in cognition and activities of daily living in Alzheimer disease, 10
Sepsis
Early goal-direct therapy reduced mortality and multiorgan dysfunction in severe sepsis or septic shock (Correction), 80
Sertraline
Sertraline maintenance treatment reduced relapse and dropouts in post-traumatic stress disorder, 69
Serum albumin
Review: Albumin increases mortality in critically ill patients, 51
Sinusitis
Adding intranasal fluticasone to cefuroxime resolved acute rhinosinusitis, 14
Smoking cessation
A preoperative smoking intervention decreased postoperative complications in elective knee or hip replacement, 7
Somatostatin
Review: Somatostatin analogues reduce blood products used in acute bleeding esophageal varices but do not reduce mortality, 93
Spinal puncture
Review: Bed rest does not prevent cervical or lumbar puncture headaches, 24
Status epilepticus
Patients who survived the first 30 days after a first episode of status epilepticus had increased mortality at 10 years, 75
Stress disorders, post-traumatic
Sertraline maintenance treatment reduced relapse and dropouts in post-traumatic stress disorder, 69
Sumatriptan
Review: Several drugs, especially triptans, are effective for pain relief in acute migraine, 103
Syncope
Cardiac pacing reduced nonaccidental falls in older adults with cardioinhibitory carotid sinus hypersensitivity, 8

T

Tennis elbow
Physiotherapy or a wait-and-see policy were the best options for lateral epicondylitis at 1 year, 65
Textbooks
Textbook descriptions of disease—where's the beef? (Editorial), A-11 (No. 1)
Thromboembolism
Review: Postmenopausal estrogen replacement therapy increases the risk for venous thromboembolism, 42
Thrombolytic therapy
Primary percutaneous intervention was more effective than thrombolytic therapy for acute myocardial infarction, 43
Tic disorders
Clonidine and methylphenidate were effective for attention deficit hyperactivity disorder in children with comorbid tics, 70
Tinea pedis
Review: Oral antifungal drugs promote cure of fungal foot infections, 97
Troponin I
An early invasive strategy was effective in the acute coronary syndrome with elevated cardiac troponin I and T levels, 1
Troponin T
An early invasive strategy was effective in the acute coronary syndrome with elevated cardiac troponin I and T levels, 1

U

Urinary plasminogen activator
Abciximab added to urokinase increased amputation-free survival in peripheral arterial occlusion of the legs, 12
Urinary tract infections
Ultrasonography plus radiography was as effective as intravenous urography for detecting abnormalities in men with proven UTI, 112
Urography
Ultrasonography plus radiography was as effective as intravenous urography for detecting abnormalities in men with proven UTI, 112

V

Vascular diseases
Review: Antiplatelet therapy prevents occlusive vascular events in high-risk patients, 5

Venlafaxine
Review: Venlafaxine is more effective than selective serotonin-reuptake inhibitors for depression, 101
Venous thrombosis
Magnetic resonance direct thrombus imaging was accurate in suspected deep venous thrombosis, 33
Review: Extended out-of-hospital prophylaxis with heparin prevents deep venous thrombosis in elective hip arthroplasty, 11
Ventricular dysfunction, left
A prophylactically implanted cardioverter defibrillator reduced all-cause mortality in myocardial infarction, 81
Vomiting
Review: Antiemetic drugs reduce nausea in early pregnancy, 67

W

Warfarin
Warfarin combined with low-dose aspirin in MI did not provide clinical benefit beyond that of aspirin alone, 47
Weight loss
Review: Advice on low-fat diets is not better than other weight-reducing diets for sustaining weight loss in obesity, 90